

Noncommutative gravity in three dimensions coupled to spinning sources

Kiyoshi Shiraishi^{1*}, Kenji Sakamoto^{2†} and Nahomi Kan^{2‡}

¹ *Faculty of Science, Yamaguchi University, Yoshida, Yamaguchi-shi, Yamaguchi 753-8512,
Japan*

² *Graduate School of Science and Engineering, Yamaguchi University, Yoshida, Yamaguchi-shi,
Yamaguchi 753-8512, Japan*

(February 1, 2008)

Abstract

Noncommutative gravity in three dimensions with vanishing cosmological constant is examined. We find a solution which describes a spacetime in the presence of a torsional source. We estimate the phase shift for each partial wave of a scalar field in the spacetime by the Born approximation.

PACS number(s): 04.60.Kz, 11.10.Kk, 11.80.Et, 11.80.Fv

Typeset using REVTeX

*Email address: shiraish@sci.yamaguchi-u.ac.jp

†Email address: b1795@sty.cc.yamaguchi-u.ac.jp

‡Email address: b1834@sty.cc.yamaguchi-u.ac.jp

I. INTRODUCTION

We expect that noncommutative field theory [1] is worth studying, because the fundamental minimal length scale in the theory may avoid singularities and infinities which arise in usual field theories. Several attempts to formulate models of gravity on noncommutative spaces have recently been developed [2–7]. For three dimensional models, a treatment based on the Chern-Simons gauge theory has been studied in [4,5,7].

In the present paper, we consider three dimensional noncommutative gravity with no cosmological term. We show an exact solution with a localized source in the theory. The size of the point-like structure is controlled by the length scale $\sqrt{\theta}$ of the theory. The investigation of scattering by the localized object reveals how we can “see” the noncommutative space by a wave probe.

The plan of the present paper is as follows: In Sec. II we give a brief description of noncommutative space for self-containedness as well as for later use of notational conventions. In Sec. III we review noncommutative gravity in three dimensions with vanishing cosmological constant. Sec. IV describes spacetime in the presence of torsional sources. The scattering of a scalar wave is studied in Sec. V. Finally, Sec. VI contains conclusion and discussion.

II. NONCOMMUTATIVE PLANE

In this section, we review two dimensional noncommutative space. For more details, please consult a comprehensive review by Douglas and Nekrasov [1].

Consider noncommutative coordinates, for example,

$$[x, y] = i\theta, \tag{1}$$

where θ is a real, positive constant. Then the “uncertainty” lies between x and y , namely,

$$\Delta x \Delta y \geq \theta, \tag{2}$$

(where a numerical factor has omitted). This implies existence of a minimal length scale $\approx \sqrt{\theta}$. If complex combinations of the coordinates, $z = x + iy$ and $\bar{z} = x - iy$, are introduced, they satisfy

$$[z, \bar{z}] = 2\theta. \quad (3)$$

There are two different representations to describe the noncommutativity; the commutative coordinate formalism with the star product and the Fock space (operator) formalism (see [1] for details). In this paper, we simply use expressions in two formalisms identically; thus we suppress the star product symbol in this paper.¹ For example, we write the equalities as

$$1 = \sum_{n=0}^{\infty} |n\rangle\langle n|, \quad z = \sqrt{2\theta} \sum_{n=0}^{\infty} \sqrt{n+1} |n\rangle\langle n+1|, \quad \bar{z} = \sqrt{2\theta} \sum_{n=0}^{\infty} \sqrt{n+1} |n+1\rangle\langle n|, \quad (4)$$

where ket and bra satisfy $z|0\rangle = 0$, $z|n\rangle = \sqrt{2\theta}\sqrt{n}|n-1\rangle$ ($n = 1, 2, \dots$), $\bar{z}|n\rangle = \sqrt{2\theta}\sqrt{n+1}|n+1\rangle$ ($n = 0, 1, 2, \dots$), and so on. Another example is

$$|n\rangle\langle n+\ell| = 2(-1)^n \sqrt{\frac{n!}{(n+\ell)!}} \left(\frac{2r^2}{\theta}\right)^{\ell/2} L_n^\ell(2r^2/\theta) e^{-r^2/\theta} e^{i\ell\varphi}, \quad (5)$$

where $r^2 = x^2 + y^2$, $z = re^{i\varphi}$ and $L_n^\ell(x)$ is the Lagurre polynomial. In particular, $|0\rangle\langle 0| = 2e^{-r^2/\theta}$.

For later use, we *define*² the inverses of z and $\bar{z}$ as

$$\frac{1}{z} \equiv \frac{1}{\sqrt{2\theta}} \sum_{n=0}^{\infty} \frac{1}{\sqrt{n+1}} |n+1\rangle\langle n| = \frac{1}{r} (1 - e^{-r^2/\theta}) e^{-i\varphi}, \quad (6)$$

$$\frac{1}{\bar{z}} \equiv \frac{1}{\sqrt{2\theta}} \sum_{n=0}^{\infty} \frac{1}{\sqrt{n+1}} |n\rangle\langle n+1| = \frac{1}{r} (1 - e^{-r^2/\theta}) e^{i\varphi}. \quad (7)$$

¹ The differentiation is expressed as $\partial_z f = [f, \bar{z}]/(2\theta)$ and $\partial_{\bar{z}} f = [z, f]/(2\theta)$. At the same time, an integration $\int d^2x f(z, \bar{z})$ in the noncommutative theory means $2\pi\theta \sum_n f_{nn}$ for $f(z, \bar{z}) = \sum_{mn} f_{mn} |m\rangle\langle n|$. [1]

²Note that here $\frac{1}{z}$ is defined in operator formalism and this differs from z^{-1} in the usual star product formalism unless $\theta = 0$.

This definition leads to $z\frac{1}{z} = \frac{1}{\bar{z}}\bar{z} = 1$, however, one can see

$$\frac{1}{z}z = \bar{z}\frac{1}{\bar{z}} = 1 - |0\rangle\langle 0|. \quad (8)$$

Thus the derivative of $\frac{1}{z}$ and $\frac{1}{\bar{z}}$ turns out to be

$$\partial_{\bar{z}}\frac{1}{z} = \frac{1}{2\theta} \left[z, \frac{1}{z} \right] = \frac{1}{2\theta} |0\rangle\langle 0| = \frac{1}{\theta} e^{-r^2/\theta}, \quad \partial_z\frac{1}{\bar{z}} = \frac{1}{2\theta} \left[\frac{1}{\bar{z}}, \bar{z} \right] = \frac{1}{2\theta} |0\rangle\langle 0| = \frac{1}{\theta} e^{-r^2/\theta}. \quad (9)$$

Interestingly enough, in the commutative limit, we find

$$\frac{1}{\theta} e^{-r^2/\theta} \xrightarrow{\theta \rightarrow 0} \pi \delta(x) \delta(y). \quad (10)$$

III. THREE DIMENSIONAL NONCOMMUTATIVE GRAVITY

Throughout this paper, we concentrate our attention on noncommutative gravity in three dimensions.³ Three dimensional Chern-Simons noncommutative gravity has recently been studied by Bañados *et al.* [4] and more recently by Cacciatori *et al.* [5]. We would like to study noncommutative gravity in three dimensions without a cosmological constant [7]. Further, we consider the case that spatial coordinates are mutually noncommutative. The signature is taken to be Euclidean, and the coordinates are denoted as

$$x^1 = x, \quad x^2 = y, \quad x^3 = \tau, \quad \text{where} \quad [x, y] = i\theta. \quad (11)$$

We define a matrix-valued dreibein one-form and a connection one-form as

$$e = e^a J_a + e^4 i, \quad \omega = \omega^a J_a + \omega^4 i, \quad (12)$$

where $e^a = e^a_\mu dx^\mu$, $\omega^a = \omega^a_\mu dx^\mu$ and J_a is expressed by using the Pauli matrix:

$$J_1 = \frac{i}{2}\sigma_1, \quad J_2 = -\frac{i}{2}\sigma_2, \quad J_3 = \frac{i}{2}\sigma_3. \quad (13)$$

A matrix-valued torsion two-form and a curvature two-form are given by: [4]

³See references [2,3] for an extension to the other dimensions.

$$\mathcal{T} = de + \omega \wedge e + e \wedge \omega, \quad \mathcal{R} = d\omega + \omega \wedge \omega. \quad (14)$$

In our notation, for example, a vacuum solution of “Einstein equation” satisfies

$$\mathcal{R} = \mathcal{T} = 0. \quad (15)$$

IV. A SOLUTION WITH A TORSIONAL SOURCE

Let us examine the following form of the dreibein:

$$e = \frac{i}{2} \left\{ d\tau + \frac{S}{2i} \left(\frac{1}{z} dz - \frac{1}{\bar{z}} d\bar{z} \right) \right\} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} + \frac{i}{2} \begin{pmatrix} 0 & dz \\ d\bar{z} & 0 \end{pmatrix}, \quad (16)$$

where S is a constant and $\omega = 0$. Note that $e^4 = \omega^4 = 0$.

Then using (14) we obtain

$$\mathcal{T} = \frac{S}{4\theta} |0\rangle\langle 0| d\bar{z} \wedge dz \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}, \quad \mathcal{R} = 0. \quad (17)$$

In the commutative limit, this corresponds to the spinning solution obtained by Deser, Jackiw and ‘t Hooft [8] with spin S and vanishing mass. For a finite θ , the torsional source has a finite extension. Note that $\text{Tr } \mathcal{T} = 0$ for any value of θ .

This one-body solution can be generalized to the N -body solution for sources located at $\mathbf{z}_a$, each carrying spin S_a , $a = 1, \dots, N$. One finds the solution as

$$e = \frac{i}{2} \left\{ d\tau + \sum_{a=1}^N \frac{S_a}{2i} \left(\frac{1}{z - z_a} dz - \frac{1}{\bar{z} - \bar{z}_a} d\bar{z} \right) \right\} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} + \frac{i}{2} \begin{pmatrix} 0 & dz \\ d\bar{z} & 0 \end{pmatrix}, \quad (18)$$

where $\frac{1}{z - z_a}$ and $\frac{1}{\bar{z} - \bar{z}_a}$ are defined by

$$\frac{1}{z - z_a} = \sum_{m=0}^{\infty} z_a^m \left(\frac{1}{z} \right)^{m+1}, \quad \frac{1}{\bar{z} - \bar{z}_a} = \sum_{m=0}^{\infty} \bar{z}_a^m \left(\frac{1}{\bar{z}} \right)^{m+1}. \quad (19)$$

V. WAVE EQUATION

Now we can write down a wave equation for a scalar field around an isolated torsional source. One of the possible expression for a scalar Φ is

$$\left(\partial_\mu e_a^\mu e_a^\nu \partial_\nu - m^2\right) \Phi = 0, \quad (20)$$

where e_a^μ is the inverse of e_μ^a . Here the determinant of the dreibein has been omitted, since it is unity for the solution in Sec. IV. Substituting the dreibein of the one-body solution in Sec. IV, we find that the equation (20) reduces to

$$\left\{-\partial_t^2 + 2(D_z D_{\bar{z}} + D_{\bar{z}} D_z) - m^2\right\} \Phi = 0, \quad (21)$$

where we have changed the signature into the Lorentzian one and used the notation

$$D_z f \equiv \partial_z f - \frac{S}{2i} \frac{1}{\mathbf{z}} \partial_t f, \quad D_{\bar{z}} f \equiv \partial_{\bar{z}} f + \frac{S}{2i} \frac{1}{\bar{\mathbf{z}}} \partial_t f. \quad (22)$$

The constant m denotes the mass of the scalar particle.

If we choose the following “monochromatic” wave with energy E as the wave function Φ :

$$\Phi(x, y, t) = \phi(z, \bar{z}) e^{-iEt}, \quad (23)$$

then the stationary function satisfies:

$$\left\{2\left(\tilde{D}_z \tilde{D}_{\bar{z}} + \tilde{D}_{\bar{z}} \tilde{D}_z\right) + k^2\right\} \phi(z, \bar{z}) = 0, \quad (24)$$

where

$$\tilde{D}_z f = \partial_z f + \frac{SE}{2} \frac{1}{\mathbf{z}} f, \quad \tilde{D}_{\bar{z}} f = \partial_{\bar{z}} f - \frac{SE}{2} \frac{1}{\bar{\mathbf{z}}} f, \quad (25)$$

and the wave number k is defined as $k = \sqrt{E^2 - m^2}$.

In the case with $S = 0$, or Minkowski space, the scalar wave equation becomes

$$\left\{2(\partial_z \partial_{\bar{z}} + \partial_{\bar{z}} \partial_z) + k^2\right\} \chi_\ell = 0, \quad (26)$$

and its regular solution χ_ℓ is expressed by using the Bessel function as

$$\begin{aligned}\chi_\ell &= J_\ell(kr)e^{i\ell\varphi} \\ &= \sum_{n=0}^{\infty} \sqrt{\frac{n!}{(n+\ell)!}} \left(\frac{k^2\theta}{2}\right)^{\ell/2} L_n^\ell(k^2\theta/2)e^{-k^2\theta/4}|n\rangle\langle n+\ell| \quad (\ell = 0, 1, 2, \dots),\end{aligned}\quad (27)$$

and its asymptotic behavior is

$$\chi_\ell \xrightarrow{r \rightarrow \infty} \sqrt{\frac{2}{\pi kr}} \cos\left(kr - \frac{\ell\pi}{2} - \frac{\pi}{4}\right) e^{i\ell\varphi}. \quad (28)$$

Returning to the $SE > 0$ case, Eq. (24) can be rewritten as

$$\left\{2\left(\tilde{D}_z\tilde{D}_{\bar{z}} + \tilde{D}_{\bar{z}}\tilde{D}_z\right) + k^2\right\}\phi_\ell = 2\left(\partial_z\partial_{\bar{z}} + \partial_{\bar{z}}\partial_z\right)\phi_\ell - U\phi_\ell + k^2\phi_\ell = 0, \quad (29)$$

where

$$U\phi_\ell = 2\alpha\left(\frac{1}{\bar{z}}\partial_z\phi_\ell - \frac{1}{z}\partial_{\bar{z}}\phi_\ell\right) + \frac{\alpha^2}{2}\left(\frac{1}{z}\frac{1}{\bar{z}} + \frac{1}{\bar{z}}\frac{1}{z}\right)\phi_\ell, \quad (30)$$

with $\alpha \equiv SE$. In this paper, we treat the case with $0 < \alpha \ll 1$ for simplicity.

Here we assume that ϕ_ℓ takes the similar form to χ_ℓ :

$$\phi_\ell = \sum_{n=0}^{\infty} C_n |n\rangle\langle n+\ell|, \quad (31)$$

where C_n depends on n , ℓ , and $k^2\theta$. This solution may have the following asymptotics:

$$\phi_\ell \xrightarrow{r \rightarrow \infty} \sqrt{\frac{2}{\pi kr}} \cos\left(kr - \frac{\ell\pi}{2} - \frac{\pi}{4} + \delta_\ell\right) e^{i\ell\varphi}, \quad (32)$$

for $r \gg k^{-1}$ and $r \gg \sqrt{\theta}$. We call a constant δ_ℓ as a phase shift. Now assuming the regularity of the solution at the origin and using asymptotic forms (28) and (32), we find

$$\int d^2x \left\{2\chi_\ell^\dagger(\partial_z\partial_{\bar{z}} + \partial_{\bar{z}}\partial_z)\phi_\ell - 2\phi_\ell^\dagger(\partial_z\partial_{\bar{z}} + \partial_{\bar{z}}\partial_z)\chi_\ell\right\} = -4\sin\delta_\ell. \quad (33)$$

Therefore the phase shift for a partial wave for $\ell \geq 0$ is written as

$$\sin\delta_\ell = -\frac{1}{4} \int d^2x \chi_\ell^\dagger U \phi_\ell. \quad (34)$$

Similarly, since $\chi_\ell^\dagger$ also is a solution for Eq. (26), we take $\chi_{-\ell} \equiv \chi_\ell^\dagger$ and $\phi_{-\ell} \equiv \phi_\ell^\dagger$ for $\ell \geq 0$.

When α is sufficiently small, the Born approximation may be valid to obtain

$$\delta_\ell \simeq -\frac{1}{4} \int d^2x \chi_\ell^\dagger U \chi_\ell, \quad \delta_{-\ell} \simeq -\frac{1}{4} \int d^2x \chi_\ell U \chi_\ell^\dagger \quad \text{for } \ell \geq 0. \quad (35)$$

The calculation done by the operator formalism leads to⁴

$$\begin{aligned} & \frac{1}{2\pi\alpha} \int d^2x \chi_\ell^\dagger (2\alpha) \left(\frac{1}{\bar{z}} \partial_z \chi_\ell - \frac{1}{z} \partial_{\bar{z}} \chi_\ell \right) \\ &= 2 \left(\frac{k^2\theta}{2} \right)^\ell e^{-k^2\theta/2} \sum_{n=0}^{\infty} \frac{n!}{(n+\ell)!} L_n^\ell(k^2\theta/2) (L_{n+1}^\ell(k^2\theta/2) - L_n^\ell(k^2\theta/2)) \\ & \quad + \frac{1}{\ell!} \left(\frac{k^2\theta}{2} \right)^\ell e^{-k^2\theta/2} (L_0^\ell(k^2\theta/2))^2 \\ &= 1 - 2 \frac{\gamma(\ell, k^2\theta/2)}{\Gamma(\ell)} + \frac{1}{\ell!} \left(\frac{k^2\theta}{2} \right)^\ell e^{-k^2\theta/2} \\ &= 1 - 2 \frac{\gamma(\ell+1, k^2\theta/2)}{\ell!} - \frac{1}{\ell!} \left(\frac{k^2\theta}{2} \right)^\ell e^{-k^2\theta/2} \quad (\ell > 0), \end{aligned} \quad (36)$$

where the incomplete gamma function

$$\gamma(\ell+1, y) = \int_0^y e^{-t} t^\ell dt, \quad (37)$$

has been used. In the same manner, one finds

$$\begin{aligned} & \frac{1}{2\pi\alpha} \int d^2x \chi_0^\dagger (2\alpha) \left(\frac{1}{\bar{z}} \partial_z \chi_0 - \frac{1}{z} \partial_{\bar{z}} \chi_0 \right) \\ &= 2e^{-k^2\theta/2} \sum_{n=0}^{\infty} L_n(k^2\theta/2) (L_{n+1}(k^2\theta/2) - L_n(k^2\theta/2)) + e^{-k^2\theta/2} (L_0(k^2\theta/2))^2 \\ &= -1 + e^{-k^2\theta/2} \quad (\ell = 0), \end{aligned} \quad (38)$$

where $L_n(x) \equiv L_n^0(x)$, as well as

$$\begin{aligned} & \frac{1}{2\pi\alpha} \int d^2x \chi_\ell (2\alpha) \left(\frac{1}{\bar{z}} \partial_z \chi_\ell^\dagger - \frac{1}{z} \partial_{\bar{z}} \chi_\ell^\dagger \right) \\ &= 2 \left(\frac{k^2\theta}{2} \right)^\ell e^{-k^2\theta/2} \sum_{n=1}^{\infty} \frac{n!}{(n+\ell)!} L_n^\ell(k^2\theta/2) (L_{n-1}^\ell(k^2\theta/2) - L_n^\ell(k^2\theta/2)) \\ & \quad - \frac{2}{\ell!} \left(\frac{k^2\theta}{2} \right)^\ell e^{-k^2\theta/2} (L_0^\ell(k^2\theta/2))^2 \\ &= -1 \quad (\ell > 0). \end{aligned} \quad (39)$$

⁴See Appendix for formulas on the Laguerre functions.

Consequently, we have the expression for the phase shift in the Born approximation up to $O(\alpha)$,

$$\delta_\ell \simeq \begin{cases} -\frac{\pi\alpha}{2} \left\{ 1 - 2\frac{\gamma(\ell+1, k^2\theta/2)}{\ell!} - \frac{1}{\ell!} \left(\frac{k^2\theta}{2}\right)^\ell e^{-k^2\theta/2} \right\} & (\ell > 0) \\ \frac{\pi\alpha}{2}(1 - e^{-k^2\theta/2}) & (\ell = 0) \\ \frac{\pi\alpha}{2} & (\ell < 0) \end{cases} \quad (40)$$

In the commutative limit, $\theta \rightarrow 0$, this result reduces to $\delta_\ell = -\pi\alpha/2$ for $\ell > 0$ and $\delta_\ell = \pi\alpha/2$ for $\ell < 0$, which coincides with the one derived by using

$$\int_0^\infty \frac{\ell}{r} (J_\ell(r))^2 dr = \frac{1}{2} \quad (\ell > 0),$$

for the ordinary integration.

In the opposite limit, $\theta \rightarrow \infty$, the phase shift for any ℓ becomes the same value

$$\delta_\ell = \frac{\pi\alpha}{2} \quad \text{when} \quad \theta \rightarrow \infty. \quad (41)$$

In the usual commutative model [9], it is known that the first-order Born approximation up to $O(\alpha)$ gives an exact result for the phase shift (except for $\ell = 0$).

VI. CONCLUSION AND DISCUSSION

To summarize, we have found an exact solution for three dimensional noncommutative gravity and studied the phase shift of scalar waves in the spinning background spacetime. The behavior of the phase shift is much alike Aharonov-Bohm scattering; especially for $\theta = 0$ case, the result of de Sousa Gerbert and Jackiw [9] is reproduced. When θ is finite, the wave of sufficiently short wave length can “see” the extension of the torsional source, then the difference in the phase shift is reduced following the order of the angular momentum ℓ .

The asymmetric result on the sign of ℓ comes from the choice of the wave equation. In noncommutative theory, left- and right- covariant derivatives give different results in general, where the right-covariant derivatives read

$$D_z^R f \equiv \partial_z f - \partial_t f \frac{S}{2i} \frac{1}{z}, \quad D_{\bar{z}}^R f \equiv \partial_{\bar{z}} f + \partial_t f \frac{S}{2i} \frac{1}{\bar{z}}, \quad (42)$$

and the previous ones (22) are the left-derivative. In other words, if Φ is a solution of the wave equation (21), $\Phi^\dagger$ is not always a solution.

The wave equation (21) can be expected to be derived from the lagrangian density

$$-(e_a^\mu \partial_\mu \Phi)^\dagger (e_a^\nu \partial_\nu \Phi) - m^2 \Phi^\dagger \Phi.$$

The lagrangian density

$$\propto -(e_a^\mu \partial_\mu \Phi)^\dagger (e_a^\nu \partial_\nu \Phi) - (\partial_\mu \Phi e_a^\mu)^\dagger (\partial_\nu \Phi e_a^\nu) - \dots$$

leads to a wave equation

$$\frac{1}{2} \partial_\mu e_a^\mu e_a^\nu \partial_\nu \Phi + \frac{1}{2} \partial_\mu \partial_\nu \Phi e_a^\nu e_a^\mu - m^2 \Phi = 0, \quad (43)$$

which includes the left- and right-covariant derivatives for our spacetime; this yields a symmetric result for the phase shift, that is

$$\delta_\ell \simeq \begin{cases} -\frac{\pi\alpha}{2} F(\ell, k^2\theta/2) & (\ell > 0) \\ 0 & (\ell = 0) \\ \frac{\pi\alpha}{2} F(|\ell|, k^2\theta/2) & (\ell < 0) \end{cases}, \quad (44)$$

with

$$F(p, q) \equiv 1 - \frac{\gamma(p+1, q)}{p!} - \frac{1}{2p!} q^p e^{-q}. \quad (45)$$

$F(p, q)$ is plotted in Fig. 1.

In this paper, we have found a noncommutative solution describing massless spinning sources. We wonder how we can obtain “conical” (*i.e.* massive) solutions in noncommutative gravity. We also want to know how we take global properties of spacetime into account. How and when do we have to use a noncommutative torus and sphere? The choice of global structure of spacetime must be a very important problem in noncommutative gravity in any dimensions and in any formulation.

ACKNOWLEDGEMENT

We would like to thank Y. Cho for reading this manuscript and for useful comments.

APPENDIX: SOME USEFUL FORMULAS FOR THE LAGUERRE FUNCTIONS

Some useful formulas for the Laguerre functions are listed as follows [10]:

$$\frac{e^{xt/(1+t)}}{(1+t)^{\ell+1}} = \sum_{n=0}^{\infty} (-1)^n L_n^{\ell}(x) t^n, \quad (\text{A1})$$

$$L_0^{\alpha}(x) = 1, \quad (\text{A2})$$

$$\sum_{n=0}^{\infty} \frac{n!}{(n+\ell)!} (-1)^n L_n^{\ell}(x) L_n^{\ell}(y) t^n = \frac{(xyt)^{-\ell/2}}{1+t} e^{t(x+y)/(1+t)} J_{\ell}(2\sqrt{xyt}/(1+t)), \quad (\text{A3})$$

$$L_n^{\alpha-1}(x) = L_n^{\alpha}(x) - L_{n-1}^{\alpha}(x), \quad (\text{A4})$$

$$\sum_{n=0}^{\infty} \frac{n!}{(n+\ell)!} L_n^{\ell}(x) L_{n+1}^{\ell-1}(x) = \frac{e^x}{2x^{\ell}} \left(1 - 2 \frac{\gamma(\ell, x)}{\Gamma(\ell)} \right), \quad (\text{A5})$$

where

$$\gamma(\ell, y) = \int_0^y e^{-t} t^{\ell-1} dt, \quad (\text{A6})$$

$$\sum_{n=0}^{\infty} \frac{n!}{(n+\ell)!} L_n^{\ell-1}(x) L_n^{\ell}(x) = \frac{e^x}{2x^{\ell}}. \quad (\text{A7})$$

REFERENCES

- [1] M. R. Douglas and N. A. Nekrasov, Rev. Mod. Phys. **73** (2001) 977, [hep-th/0106048](#).
- [2] V. P. Nair, [hep-th/0112114](#).
- [3] A. H. Chamseddine, [hep-th/0202137](#).
- [4] M. Bañados, O. Chandía, N. Grandi, F. A. Shaposnik and G. A. Silva, Phys. Rev. **D64** (2001) 084012.
- [5] S. Cacciatori, D. Klemm, L. Martucci and D. Zanon, [hep-th/0201103](#).
- [6] S. Cacciatori, A. H. Chamseddine, D. Klemm, L. Martucci, W. A. Sarba and D. Zanon, [hep-th/0203038](#).
- [7] K. Shiraishi, [hep-th/0202099](#).
- [8] S. Deser, R. Jackiw and G. 't Hooft, Annals of Physics (NY) **152** (1984) 220.
- [9] P. de Sousa Gerbert and R. Jackiw, Commun. Math. Phys. **124** (1989) 229.
- [10] I. S. Gradstein and I. M. Ryshik, *Tables of integrals, sums, series and products*, Nauka, Moscow (1971).

FIGURES




FIG. 1. $F(p, q)$ is plotted against q . Each line is drawn for $p = 1, 2, 3, \dots$, from the left to the right.